
Project Stats

- 63.5 acres
- 84 parcels purchased from 60 property owners over 15 year period via willing-seller program
- Removed 50,000 cubic vards of soil and other material
- Created approximately 140 acre-feet of flood storage (up to about 6-year storm event)
- Removed 845 trees, including many invasive trees; salvaged timber was reused onsite as instream large woody debris, brush piles, or wood chips
- Enhanced 3,800 linear feet of streambank
- Removed three bridges and roads
- Installed public access amenities, including multi-modal trail and parking lot

Constructed in 2011 (phase 1)



Special Status Fish Species Salvaged Before Construction

- Coho salmon
- Cutthroat trout
- Steelhead
- Lamprey









Floodplain Restoration: Reducing flooding, improving water quality, and enhancing wildlife habitat in Johnson Creek www.portlandoregon.gov bes/fosterfloodplain

Foster Floodplain Natural Area Restoration Walk

> Society for Ecological Restoration, NW Chapte chapter.ser.org/northwest

> > Saturday August 24th, 2013 9:00 a.m. to noon

Stormwater runoff can impact water quality in rivers and streams. Natural resources, like urban streams, forests, and wetlands manage stormwater naturally to protect water quality, public health, and the environment. Green infrastructure reduces flooding and erosion, provides cool, clean water to our rivers and streams, and adds wildlife habitat and urban green space for healthier watersheds.



Healthy Parks, Healthy Portland






Native Plant Communities

- Riparian forest
 - 20.2 acres new
 - 9.7 acres enhanced
- Upland forest
 - 4.1 acres new
 - 4.3 acres enhanced
- Scrub-shrub and forested wetland
 - 16.8 acres new
 - 1.3 acres enhanced
 - Existing willow scrub-shrub and Oregon ash—slough sedge wetlands on site
- Other: Oak woodland (0.5 acre), emergent and herbaceous wetlands, water quality facilities, street trees, parking lot landscaping

Target Invasive Plants

- Garlic mustard (Alliaria petiolata)
- Himalayan blackberry (Rubus armeniacus)
- English ivy (Hedera helix)
- Japanese knotweed (Polygonum cuspidatum)
- Sycamore maple (Acer pseudoplatanus)
- English hawthorn (Crataegus monogyna)
- Sweet cherry (Prunus avium)
- Norway maple (Acer platanoides)
- Black locust (Robinia pseudoacacia)



FOSTER FLOODPLAIN NATURAL AREA—1998



FOSTER FLOODPLAIN NATURAL AREA—2012

